

Base map derived from U.S. Geological Survey (USGS) 7.5-minute Series (Ctopographic).  
 Woodsboro Quadrangle 1953 (Photorevised 1986).  
 Hydrology (Hydro) shown as from USGS digital line graphs (DLG) for this quadrangle.  
 Topography and cultural transportation layers from USGS stable-base film separates.  
 (Topography by photogrammetric methods from aerial photographs taken 1943).  
 Culture revised by USGS 1953.  
 Map dated in 1986 by USGS based on aerial photographs taken 1982 and other sources.  
 This information not field checked and may not meet USGS content standards.

1986 magnetic north declination (center of quadrangle): 9.5 degrees west  
 (To determine current magnetic declination see: <http://www.ngdc.noaa.gov/cgi-bin/seg/gmap/10dnh1.pl>)

Current map projection:  
 Maryland State Plane Coordinate System 1987  
 (Projection: Lambert Conformal Conic, 1980 geoidic reference system)  
 (Horizontal Datum: North American Datum 1983)  
 MD State Plane 2000-meter grid ties and coordinates shown in black.  
 Geographic coordinates (latitude/longitude) shown near corners and 2.5' intervals (in black).

Description of Map Units

- Quaternary**
  - Alluvium**  
Well to poorly sorted, stratified mixtures of unconsolidated clay, silt, sand, gravel, and cobbles; underlies flood plains of nearly all rivers and tributaries. The channel of the tributary is commonly on bedrock with alluvium exposed along the banks. Thickness of alluvium is highly variable as a function of bedrock topography, and landscape practices. Thickness estimated at as much as 15 feet (4 to 5 m) along the Monocacy River.
  - Colluvium**  
Unsorted, light gray to reddish gray, angular boulders to cobbles of quartzite and vein quartz with a silty, clayey reddish matrix. Thickness ranges from a thin veneer to more than 100 feet (30 m).
  - Weathering residuum**  
Mixture of moderate reddish brown soil and pebbles to blocks of grayish pink to white angular locally euhedral quartz. Thickness ranges from a thin veneer to 10 feet (3.0 m) thick.
  - Terrace deposits**  
Reddish brown to brown, sandy and clayey mixture of rounded pebbles to cobbles of sandstone, vein quartz and quartzite. Present along elevated low relief areas above the Monocacy River. Several separate levels of terrace deposits can be observed, but are not separately mapped here. Thickness ranges from a thin veneer to more than 10 feet (3 m).
- Jurassic**
  - Diabase dike(s)**  
Medium to dark gray, medium crystalline and equigranular, massive dikes, with characteristic orange-brown weathered surface. Dikes range from a wedge-edge to more than 150 feet (46 m) wide. While displayed as a solid unit on the map, the area mapped as diabase dike(s) actually represents an interfingering with the surrounding bedrock unit(s).
- Triassic**
  - Gettysburg Formation**  
Reddish brown to reddish gray, locally greenish gray, medium-grained siltstone, variegated claystone and fine-grained sandstone. Sandstones are thin and discontinuous. Siltstone units are root mottled and contain light gray calcitic carbonate nodules. The thickness of this member is estimated at 8,000 feet (2,440 m) (Brezinski, 2004).
  - New Oxford Formation**  
Interbedded, light reddish gray to reddish brown, root-mottled mudstone, reddish brown sandstone, and reddish siltstone. Most sandstone units are medium to coarse grained, arkosic, have basal lag conglomerates, and fine up-section. A few are relatively laterally continuous and mappable (fms). Mudstone intervals contain pervasive root casts, and local calcitic intervals. A brownish red to reddish gray, limestone to quartz-pebble conglomerate (limestone) that is 30 to 60 feet (9 to 18 m) thick, marks the base of the formation. The formation may be as thick as 10,000 feet (3,048 m) (Brezinski, 2004).
  - Grove Formation**  
Thick-bedded to massive, medium to light gray limestone with interbedded tan to medium gray dolomite. These members are recognized and mapped in the Frederick Valley as identified by Brezinski (2004).
  - Woodsboro Member**  
Interbedded, thin-bedded, dark gray, fine-grained limestone, medium-bedded, bioturbated, dark gray, fine-grained limestone, and thin wavy bedded limestone with tan, dolomitic partings. Top of Woodsboro Member not exposed because it is covered by the Triassic New Oxford Formation. Thickness is 500 feet (152 m).
  - Fountain Rock Member**  
Very thick-bedded, medium light gray, locally sandy, with thick intervals of massive, algal thrombolites, and medium-bedded, cross-bedded, sandy limestone. Two cross-bedded sandstone units recognized in the middle of the formation (Oy6) were identified and traced. Thickness is in question, but may be as much as 3,300 feet (1,006 m) along the western flank of the syncline in the Woodsboro Quadrangle, and as little as 1,000 feet (305 m) on the eastern flank.
  - Ceresville Member**  
Medium light gray to medium gray, thick-bedded and cross-bedded, arenaceous limestone and sandy dolomitic limestone with thin interbeds (1 ft or 0.3 m) of medium light gray, sandy thrombolitic dolomite. Thickness is approximately 150 to 200 feet (46 to 61 m).
- Ordovician**
  - Frederick Formation**  
Very thin to medium-bedded limestone, dolomite and thin intervals of shale, sandstone and breccia. Four members are recognized and mapped within the Frederick Formation and were named by Reinhardt (1974) and Brezinski (2004).
  - Lime Kill Member**  
Interbedded, thinly laminated to thinly bedded, dark gray, fine-grained limestone, calcareous shale, and medium-bedded, fine-grained limestone near the base, becoming more thickly interbedded toward the top with medium dark gray, fine-grained limestone and wavy bedding and stromatolitic algal beds. Near the top, the member becomes interbedded with cross-bedded, sandy, medium light gray limestone. Thickness is 700 feet (213 m).
- Cambrian**
  - Adamstown Member**  
Thinly interbedded, medium dark gray to dark gray, argillaceous, fine-grained limestone and dusky yellow to medium dark gray, silty dolomite. Limestone beds range from 0.1 to 1.5 inches (0.2 to 4.0 cm) in thickness. Several thin (6 to 16 ft, or 2 to 5 m), dark greenish gray to greenish black, light olive brown weathering, silty, calcareous shale intervals are present throughout the member. The top of the member is capped at the base of the lowest medium to thick bed of sandy or algal limestone. Thickness is approximately at 1000 feet (303 m).
  - Rocky Springs Station Member**  
Nodular to lumpy-bedded, dark gray, argillaceous, dolomitic limestone at the base with an interval of grayish black, platy shale. Bedding becoming laminated to flaggy up-section. The dark gray limestone with dusky yellow to light olive gray limestone with silty, dolomitic partings and laminations is interbedded with intervals (1 to 33 ft, or 0.3 to 10 m) of thick-bedded, medium dark gray, polymictic breccia that grade up-section into planar-bedded, arenaceous, medium gray limestone. Clast sizes range from sand size to 1 foot (0.3 m) on the diameter on the western flank of the Frederick Valley syncline and diminish to less than 1 to 2 inches (3 to 5 cm) in diameter on the eastern flank. Top of the member is mapped at the top of the stratigraphically highest polymictic breccia or sandstone interval. Thickness is approximately at 1,200 feet (366 m) in the Woodsboro Quadrangle.
  - Monocacy Member**  
Black platy shale at the base and top, with polymictic breccia beds, and thinly laminated, shaly limestone in the middle. A thick interval of black shale that marks the top of the member (Cm5) was separately mapped. Thickness is approximately 200 feet (60 m).
  - Araby Formation**  
Thickly bedded, greenish black to grayish black, very fine grained to fine-grained, burrow-mottled, silty sandstone, interbedded with medium gray to grayish black, phyllitic shales 3 to 10 feet (1 to 3 m) thick. Top of the formation consists of grayish black, phyllitic shale 50 to 65 feet (15 to 20 m) thick. The Araby Formation is present on the eastern side of the Frederick Valley Syncline, and is considered correlative to the Annetam Sandstone on the western side of the syncline. Thickness of the Araby Formation is estimated at 600 feet (183 m).
  - Sams Creek Formation**  
Dark greenish gray to medium bluish gray, argillaceous to porphyritic, massive to schistose metasediments composed of chlorite, epidote, quartz, altered plagioclase, actinolite, hornblende, and albite. Igneous texture is locally preserved and pods of epidote are common; includes some metaconglomerate composed of greenstone pebbles and cobbles, and local pillow structures and hydroclastic.
  - Marble**  
Interbedded, light olive-gray to olive-gray, medium- to coarse-grained, medium-bedded, locally ferruginous, micaceous, silty sandstone and very fine grained, silty sandstone to sandy siltstone. Thickness is estimated at 300 feet (100 m).
  - Tuffaceous phyllite**  
Grayish red-purple and bluish gray, variegated, vesicular phyllite with light gray streaks and blebs of tuffaceous phyllite.
  - Quartzite**  
Light gray to grayish green, medium-grained, thin-bedded to massive quartzite and minor calcareous sandstone. Contains detrital plagioclase, orthoclase, and polymictic quartz. Bedding is defined by concentrations of heavy minerals. Light gray, medium- and coarse-grained quartzite, locally calcareous and cross-bedded also present.
  - Ipswich Phyllite**  
Dusky blue, grayish blue, very dark red-purple, greenish gray to pale olive phyllite. Phyllonite contains abundant pods and folded stringers of white vein quartz, and minor silt. Intensely folded and sheared with finely laminated beds seen only in slate. Phyllite consists mostly of muscovite and chlorite, but also contains paragonite and chloritoid. Has a texture sheen because of paragonite (determined by x-ray diffraction) and dark color because of abundant hematite dust.
  - Conglomeratic metagraywacke**  
Light gray and green, medium- to coarse-grained metagraywacke with white quartz pebbles, variegated phyllite, and green chloritic matrix.
  - Chlorite phyllite**  
Light olive-gray and greenish gray, chlorite phyllite and metasilstone.

References

Brezinski, D.K., 2004. Stratigraphy of Frederick Valley and its relationship to karst development. Maryland Geological Survey, Report of Investigations 75, 101 p.

Brezinski, D.K. and Edwards, J., Jr., 2004. Geologic map of the Woodsboro Quadrangle, Frederick and Carroll Counties, Maryland. Maryland Geological Survey, Quadrangle Map, scale 1:24,000 (Version WOODSGE02004.1).

Reinhardt, J., 1974. Stratigraphy, Sedimentology and Cambro-Ordovician Paleogeography of the Frederick Valley, Maryland: Maryland Geological Survey, Report of Investigations 23, 73 p.

Supplemental Information

**Use Constraints:** These data represent the results of data collection/processing for a specific Department of Natural Resources, Maryland Geological Survey activity and indicate general existing conditions. As such, they are only valid for the intended use, content, time, and accuracy specifications. The user is responsible for the results of any application of the data for other than their intended purpose. The Maryland Geological Survey makes no warranty, expressed or implied, as to the use or appropriateness of the data, and there are no assurances of merchantability or fitness for a particular purpose of use. The Maryland Geological Survey makes no representation as to the accuracy or completeness of the data and may not be held liable for human error or defect. Data are only valid at 1:24,000 scale. Data should not be used at a scale greater than that.

**Acknowledgements:** This map was funded in part by the Maryland State Highway Administration.

Field mapping of karst features was conducted by David Brezinski in 2002-2004. Field mapping of the geology was completed in 2003-2004. This karst map was compiled in digital form by Heather Quinn of the Maryland Geological Survey and Brent Anderson and Catherine Lusksholtz of Towson University, Center for Geographic Information Sciences.

The facilities and services of the Maryland Department of Natural Resources are available to all without regard to race, color, religion, sex, sexual orientation, age, national origin or physical or mental disability.

Version: WOODSGE02004.1  
 Released June 2004

STATE OF MARYLAND  
 Robert E. Ehrlich, Jr.  
 Governor  
 Michael S. Steele  
 Lieutenant Governor



DEPARTMENT OF NATURAL RESOURCES  
 C. Ronald Franks  
 Secretary


W. P. Jensen  
 Deputy Secretary  
 MARYLAND GEOLOGICAL SURVEY  
 Emery T. Cleaves  
 Director

Copies of this map are available in hard copy (paper) and digital form from: MARYLAND GEOLOGICAL SURVEY, 2300 Saint Paul Street, Baltimore, MD 21218. Ph: 410-554-5500, Fax: 410-554-5500, <http://www.mgs.net/gv/>



Karst Features of the Woodsboro Quadrangle, Frederick and Carroll Counties, Maryland

By David K. Brezinski 2004

Scale 1:24,000



Contour Interval 20 Feet  
 National Geodetic Vertical Datum of 1929  
 (To convert elevations to the North American Vertical Datum of 1988, subtract 1 foot)  
 (To convert from feet to meters, multiply by 0.3048)



Adjoining 7.5' Quadrangle Names  
 Woodsboro Quadrangle, shaded

1	2	3
4	5	6
7	8	

1. Blue Ridge Summit  
 2. Emmittsburg  
 3. Taneytown  
 4. Carolina Furnace  
 5. Union Bridge  
 6. Frederick  
 7. Walkersville  
 8. Libertytown

**Explanation of Map Symbols**

**Geologic Symbols**

**Contacts**  
 Geologic contact; approximately located  
 dotted where concealed

**Karst Features**  
 Active Sinkhole  
 Depression  
 Spring

**Faults**  
 U Uplifted side  
 D Downthrown side  
 Thin fault  
 sawtooth on upthrown block  
 Overturned Thrust Fault  
 base of sawtooth on upper plate;  
 sawtooth in direction of dip  
 Fault; concealed  
 Fault; inferred

**Base Map Symbols**

**Topographic and Hydrologic Symbols**  
 Topographic Contour - Index (100-ft interval)  
 Topographic Contour - Intermediate (20-ft interval)  
 Stream  
 Water body (including lakes, ponds, streams)

For additional information on USGS cultural/transportation symbology see <http://pubs.usgs.gov/of/1999/of99-430/>