

- (vii) Streptococcus meningitis type A.
 - (viii) Streptococcus meningitis type B.
 - (ix) Viral meningitis.
 - (x) Meningococemia.
 - (xi) Typhoid or nontyphoid salmonellosis.
 - (xii) Syphilis.
 - (xiii) Tuberculosis.
 - (xiv) Human immunodeficiency virus infection.
 - (xv) CD 4+ count, if less than 200/MM3.
 - (xvi) Chlamydia infection.
 - (xvii) Crypto sporidiosis.
 - (xviii) E. coli 0157:H7 infection.
 - (xix) Lyme disease.
 - (xx) Pertussis.
 - (xxi) Rocky Mountain spotted fever.
 - (xxii) Shigellosis.]
- (1) AMOEBIASIS.
 - (2) ANTHRAX.
 - (3) BACTEREMIA IN NEWBORNS.
 - (4) BOTULISM.
 - (5) BRUCELLOSIS.
 - (6) CAMPYLOBACTER INFECTION.
 - (7) CD 4+ COUNT, IF LESS THAN 200/MM3.
 - (8) CHLAMYDIA INFECTION.
 - (9) CHOLERA.
 - (10) COCCIDIOIDOMYCOSIS.
 - (11) CRYPTOSPORIDIOSIS.
 - (12) CYCLOSPORIASIS.
 - (13) DENGUE FEVER.